

PHARMACOPŒIA
OF THE
ROYAL LONDON
OPHTHALMIC HOSPITAL

K50271



22500289336

Med

K50271

PHARMACOPŒIA.

LONDON : PRINTED BY
SPOTTISWOODE AND CO., NEW-STREET SQUARE
AND PARLIAMENT STREET

85770

THE
PHARMACOPŒIA
OF THE
ROYAL LONDON
OPHTHALMIC HOSPITAL.

SEVENTH EDITION.

LONDON : 1882.

17 694 788

WELLCOME INSTITUTE LIBRARY	
Coll.	WelMOmec
Coll.	
No.	WW

Consulting Physician.

~~FREDERICK JOHN FARRE, M.D.~~

Physician.

~~ROBERT MARTIN, M.D.~~

Consulting Surgeons.

~~JAMES DIXON.~~

~~GEORGE CRITCHETT.~~

~~WILLIAM BOWMAN, F.R.S.~~

~~JONATHAN HUTCHINSON.~~

Surgeons.

~~JOHN CAWOOD WORDSWORTH.~~

~~JOHN FREMLYN STREETFELD.~~

~~JOHN WHITAKER HULKE, F.R.S.~~

~~GEORGE LAWSON.~~

JOHN COUPER.

Assistant Surgeons.

WARREN TAY.

~~JAMES EDWARD ADAMS.~~

JOHN TWEEDY.

~~ROBERT LYELL.~~

PHARMACOPŒIA.



CAUSTICA — CAUSTICS.

CAUSTICUM ZINCI CHLORIDI.

(MIDDLESEX HOSPITAL.)

Take of

Chloride of Zinc	.	.	480 grains
Wheat Flour	.	.	180 grains
Water	.	.	1 fluid ounce

Mix, and apply the heat of a water bath
for a *few minutes only*, until a paste is
formed.

Or—

Take of

Chloride of Zinc	.	.	} equal parts
Oxide of Zinc	.	.	

Mix thoroughly with pestle and mortar,
and preserve in a stoppered bottle.

*CONFECTIO — CONFECTION.***CONFECTIO SULPHURIS SIMPLEX.**

Take of

Sublimed Sulphur	.	.	4 ounces
Acid Tartrate of Potash	.	.	1 ounce
Treacle, by weight	.	.	4 ounces

Rub well together.

Dose, one or two teaspoonfuls for an adult.

*DECOCTA — DECOCTIONS.***DECOCTUM PAPAVERIS.**

Take of

Poppy Capsules, bruised (without seeds)	.	.	1 ounce
Distilled Water	.	.	1 pint

Boil for a quarter of an hour in a covered vessel, and strain.

DECOCTUM QUERCUS.

Take of

Oak Bark, bruised	.	.	1 ounce
Distilled Water	.	.	1 pint

Boil for a quarter of an hour in a covered vessel, and strain.

FOMENTUM — FOMENTATION.

FOMENTUM BELLADONNÆ.

Take of

Extract of Belladonna	. 60 grains
Distilled Water 1 pint

Rub together till dissolved.

To be made hot and used as a fomentation.

GARGARISMATA — GARGLES.

GARGARISMA ACIDI CARBOLICI.

Take of

Glycerine of Carbolic Acid	
B.P. 2½ fluid drachms
Glycerine 3 fluid drachms
Water up to 20 fluid ounces

Mix.

GARGARISMA ACIDI TANNICI.

Take of

Diluted Hydrochloric Acid	2 fluid drachms
Glycerine of Tannin B.P. .	3 fluid drachms
Water	up to 20 fluid ounces

Mix.

GARGARISMA ALUMINIS.

Take of

Powdered Alum	.	.	180 grains
Glycerine	.	.	1 fluid ounce
Distilled Water	.	up to	20 fluid ounces

Dissolve and mix.

GARGARISMA HYDRARGYRI PERCHLORIDI.

Take of

Perchloride of Mercury	.	2 grains
Tincture of Cochineal	.	2 fluid drachms
Water	.	up to 20 fluid ounces

Dissolve and mix.

GARGARISMA POTASSÆ CHLORATIS.

Take of

Chlorate of Potash	.	.	90 grains
Hydrochloric Acid	.	.	40 minims
Glycerine	.	.	1 fluid ounce
Distilled Water	.	up to	20 fluid ounces

Mix the acid and chlorate in a bottle. Cork and let it stand for three minutes. Then add the water gradually with much shaking. Lastly add the glycerine.

GARGARISMA SODÆ CHLORATÆ.

Take of

Solution of Chlorinated	
Soda	1 ounce
Distilled Water	19 ounces

Mix.

GLYCERINA — GLYCERINES.

GLYCERINUM ACIDI TANNICI P.B.

Take of

Tannic Acid	1 ounce
Glycerine	4 fluid ounces

Dissolve with gentle heat.

GLYCERINUM BORACIS P.B.

Take of

Borax, in powder	1 ounce
Glycerine	4 fluid ounces

Rub together till dissolved.

GLYCERINUM BELLADONNÆ.

Take of

Extract of Belladonna	. . .	1 ounce
Distilled Water	4 fluid drachms
Glycerine	3½ fluid ounces

Mix the extract with the water, add the glycerine, and rub together till dissolved.

GLYCERINUM PLUMBI SUBACETATIS.

Take of

Acetate of Lead	5 ounces
Oxide of Lead, in powder	3½ ounces
Glycerine	20 fluid ounces
Distilled Water	12 fluid ounces

Mix together and boil for a quarter of an hour; then filter and evaporate to one pint.

*GUTTÆ — DROPS.***GUTTÆ ARGENTI NITRATIS.**

Take of

Nitrate of Silver	1 grain
Distilled Water	1 fluid ounce

Dissolve.

GUTTÆ ARGENTI NITRATIS FORTIORES.

Take of

Nitrate of Silver	.	.	2 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

GUTTÆ ATROPIÆ SULPHATIS.

Take of

Sulphate of Atropia	.	.	1 grain
Distilled Water	.	.	1 fluid ounce

Dissolve.

GUTTÆ ATROPIÆ SULPHATIS FORTIORES.

Take of

Sulphate of Atropia	.	.	2 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

GUTTÆ ATROPIÆ CUM ALUMINE.

Take of

Sulphate of Atropia	.	.	1 grain
Alum	.	.	3 grains
Distilled Water.	.	.	1 fluid ounce

Dissolve.

GUTTÆ ATROPIÆ CUM ZINCO.

Take of

Sulphate of Atropia	.	.	1 grain
Sulphate of Zinc	.	.	1 grain
Distilled Water	.	.	1 fluid ounce

Dissolve.

GUTTÆ CUPRI SULPHATIS.

Take of

Sulphate of Copper	.	.	2 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

GUTTÆ DATURIÆ.

Take of

Sulphate of Daturine	.	.	2 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

GUTTÆ ESERIÆ.

Take of

Sulphate of Eserine	.	.	2 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

GUTTÆ OPII.

Take of

Wine of Opium	.	.	2 fluid drachms
Distilled Water	.	.	6 fluid drachms

Mix.

GUTTÆ PILOCARPIÆ.

Take of

Nitrate of Pilocarpine	.	.	2 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

GUTTÆ POTASSII IODIDI.

Take of

Iodide of Potassium	.	.	4 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

GUTTÆ TEREBINTHINÆ.

Take of

Oil of Turpentine	.	.	1 fluid drachm
Olive Oil	.	.	7 fluid drachms

Mix.

GUTTÆ ZINCI CHLORIDI.

Take of

Chloride of Zinc	.	.	1 grain
Distilled Water	.	.	1 fluid ounce

Dissolve.

GUTTÆ ZINCI SULPHATIS.

Take of

Sulphate of Zinc	.	.	2 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

GUTTÆ ZINCI SULPHATIS ET ALUMINIS.

Take of

Sulphate of Zinc	.	.	1 grain
Alum	.	.	3 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

INJECTIONES — INJECTIONS.

INJECTIO ALKALINA.

Take of

Carbonate of Potash	.	.	120 grains
Distilled Water	.	.	1 pint

Dissolve.

INJECTIO ASTRINGENS.

Take of

Alum in powder	.	.	80 grains
Sulphate of Zinc	.	.	80 grains
Tannic Acid	.	.	30 grains
Distilled Water	.	.	1 pint

Dissolve.

INJECTIO PLUMBI.

Take of

Diluted Solution of Sub-			
acetate of Lead	.	.	1½ fluid drachm
Rectified Spirit	.	.	2 fluid drachms
Distilled Water	.	up to	1 pint

Mix.

INJECTIONES HYPODERMICÆ — HYPO- DERMIC INJECTIONS.

INJECTIO ATROPIÆ SULPHATIS.

Take of

Sulphate of Atropia	.	.	¼ grain
Distilled Water	.	.	1 fluid drachm

Mix.

Dose, by subcutaneous injection, two to four minims. (Three minims contain $\frac{1}{80}$ th of a grain of the sulphate of atropia.)

INJECTIO MORPHIÆ HYPODERMICA P.B.

A Solution of Acetate of Morphia containing one grain of acetate in twelve minims of the injection.

Dose, one to six minims.

INJECTIO MORPHIÆ ET ATROPIÆ.

Take of

Sulphate of Atropia . . .	1 grain
Hypodermic Injection of Morphia B.P. . . .	6 fluid drachms

Mix.

Dose, one to four minims. Three minims contain $\frac{1}{4}$ th of a grain of acetate of morphia and $\frac{1}{120}$ th of sulphate of atropia.

LINCTUS.

Take of

Treacle	2 fluid ounces
Powdered Tragacanth . . .	40 grains
Tincture of Opium	30 minims
Water	2 fluid ounces
Diluted Sulphuric Acid . . .	30 minims
Vinegar of Squills	3 fluid drachms

Mix.

Dose, one or two teaspoonfuls.

LINIMENTA — LINIMENTS.

LINIMENTUM ACONITI ET SAPONIS.

Take of

Liniment of Aconite B.P	1 part
Liniment of Soap . . .	3 parts

Mix.

LINIMENTUM AMMONIÆ P.B.

Take of

Solution of Ammonia . . .	1 part
Olive Oil	3 parts

Mix.

**LINIMENTUM BELLADONNÆ COM-
POSITUM.**

Take of

Extract of Belladonna . . .	1 fluid drachm
Distilled Water	1 fluid drachm
Liniment of Soap	6 fluid drachms

Mix.

LINIMENTUM CROTONIS

Take of

Croton Oil	1 fluid drachm
Oil of Turpentine	3 fluid drachms
Olive Oil	4 fluid drachms

Mix.

LINIMENTUM HYDRARGYRI OLEATIS

(10 per cent.)

(UNIVERSITY COLLEGE HOSPITAL.)

Take of

Yellow Oxide of Mercury	1 drachm
Oleic Acid	. / . . 10 fluid drachms

To the oleic acid, kept agitated in a mortar, sprinkle in the oxide of mercury gradually; triturate the mixture frequently during twenty-four hours until the oxide is dissolved and a viscid solution is formed.

LINIMENTUM HYDRARGYRI OLEATIS CUM MORPHIA.

(UNIVERSITY COLLEGE HOSPITAL.)

Take of

Morphia (pure alkaloid)	. 10 grains
Oleic Acid	. . . 5 fluid drachms

Dissolve and add

Liniment of Oleate of Mercury (10 per cent.)	. 5 drachms
--	-------------

Mix.

LIQUORES — SOLUTIONS.

LIQUOR ARGENTI NITRATIS

Take of

Nitrate of Silver	.	.	5, 10, 15, or 20
			grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

N.B.—These preparations are intended only for application to the surface of the conjunctiva of the lids with a camel's-hair brush, the excess to be afterwards neutralised with a solution of salt and water five grains to the ounce.

LIQUOR ATROPIÆ SULPHATIS P.B.

Take of

Sulphate of Atropia	.	.	4 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

LIQUOR EPISPASTICUS.

Take of

Cantharides, in powder	.	8 ounces
Glacial Acetic Acid	.	2 fluid ounces
Ether	.	a sufficiency

Mix the cantharides with the acid and two fluid ounces of ether; pack in a percolator, and after twenty-four hours pass ether slowly through until twenty fluid ounces are obtained. Keep in a stoppered bottle.

LOTIONES — LOTIONS.

LOTIO ACETI.

Take of

Diluted Acetic Acid	.	30 minims
Distilled Water	.	up to 1 fluid ounce

Mix.

LOTIO ACIDI HYDROCYANICI.

Take of

Diluted Hydrocyanic Acid	5 minims
Distilled Water	1 fluid ounce

Mix.

LOTIO ACIDI HYDROCYANICI CUM SODA.

Take of

Bicarbonate of Soda . . .	5 grains
Borax	5 grains
Diluted Hydrocyanic Acid	5 minims
Distilled Water	1 fluid ounce

Dissolve and mix.

LOTIO ALUMINIS.

Take of

Alum, in powder	6 grains
Distilled Water	1 fluid ounce

Dissolve.

LOTIO ALUMINIS MITIS.

Take of

Alum, in powder	3 grains
Distilled Water	1 fluid ounce

Dissolve.

LOTIO ALUMINIS COMPOSITA.

Take of

Alum, in powder	3 grains
Sulphate of Zinc	1 grain
Distilled Water	1 fluid ounce

Dissolve.

LOTIO ARNICÆ.

Take of

Tincture of Arnica	.	.	20 minims
Distilled Water	.	.	1 fluid ounce

Mix.

LOTIO BELLADONNÆ CUM ALUMINE.

Take of

Extract of Belladonna	.	.	5 grains
Alum, in powder	.	.	3 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

LOTIO BORACIS.

Take of

Glycerine of Borax	.	.	1 fluid drachm
Distilled Water	.	.	1 fluid ounce

Dissolve.

LOTIO FERRI SULPHATIS.

Take of

Sulphate of Iron, in powder	.	.	5 grains
Distilled Water	.	.	1 fluid ounce

Dissolve.

LOTIO HYDRARGYRI PERCHLORIDI.

Take of

Perchloride of Mercury	.	$\frac{1}{8}$ grain
Distilled Water	.	1 fluid ounce

Dissolve.

LOTIO PLUMBI ACETATIS.

Take of

Acetate of Lead, in powder	2 grains
Diluted Acetic Acid	2 minims
Distilled Water	1 fluid ounce

Dissolve and mix.

LOTIO PLUMBI FORTIS.

Take of

Solution of Subacetate of	
Lead	16 minims
Rectified Spirit	16 minims
Distilled Water	up to 1 fluid ounce

Mix.

LOTIO POTASSÆ CHLORATIS.

Take of

Chlorate of Potash	.	.	5 grains
Diluted Hydrochloric Acid	.	.	2 minims
Distilled Water	.	.	1 fluid ounce

Dissolve and mix.

LOTIO RUBRA.

Take of

Sulphate of Zinc	.	.	1 grain
Compound Tincture	of		
Lavender	.	.	15 minims
Spirit of Rosemary	.	.	15 minims
Distilled Water	.	up to	1 fluid ounce

Dissolve and mix.

LOTIO ZINCI SULPHATIS.

Take of

Sulphate of Zinc	.	.	1 grain
Distilled Water	.	.	1 fluid ounce

Dissolve.

*MISTURÆ—MIXTURES.*¹

MISTURA ACIDI HYDROCHLORICI.

Take of

Diluted Nitro-hydrochloric	
Acid	10 minims
Tincture of Chiretta	15 minims
Water ,	up to 1 fluid ounce

Mix.

**MISTURA ACIDI SULPHURICI CUM
MAGNESIA.**

Take of

Sulphate of Magnesia	60 grains
Diluted Sulphuric Acid	10 minims
Glycerine	30 minims
Peppermint Water	up to 1 fluid ounce

Mix.

MISTURA ÆTHERIS COMPOSITA.

Take of

Spirit of Ether	30 minims
Aromatic Spirit of Am-	
monia	30 minims
Syrup of Tolu	1 fluid drachm
Distilled Water	1 fluid ounce

Mix.

¹ The ordinary dose of all the mixtures is one ounce for an adult.

MISTURA AMMONIÆ ACETATIS.

Take of

Solution of Acetate of Am-	
monia	2 fluid drachms
Distilled Water	up to 1 fluid ounce

Mix.

MISTURA AMMONIÆ ACETATIS CUM
MAGNESIA

(VEL MISTURA SALINA CATHARTICA).

Take of

Sulphate of Magnesia	60 grains
Mixture of Acetate of Am-	
monia	1 fluid ounce

Mix.

MISTURA AMMONIÆ CUM IPECACUANHA.

Take of

Carbonate of Ammonia	3 grains
Wine of Ipecacuanha	10 minims
Glycerine	30 minims
Distilled Water	up to 1 fluid ounce

Dissolve and mix.

MISTURA BISMUTHI.

Take of

Subnitrate of Bismuth	.	10 grains
Powdered Tragacanth	.	3 grains
Distilled Water	.	1 fluid ounce

Mix.

MISTURA CHLOROFORMI CUM AMMONIA.

Take of

Carbonate of Ammonia	.	3 grains
Spirit of Chloroform	.	15 minims
Infusion of Gentian	.	up to 1 fluid ounce

Dissolve and mix.

MISTURA COLCHICI.

Take of

Wine of Colchicum	.	20 minims
Carbonate of Magnesia	.	10 grains
Pimento Water	.	up to 1 fluid ounce

Mix.

MISTURA COPAIBÆ.

Take of

Balsam of Copaiva	.	20 minims
Solution of Potash	.	10 minims
Pimento Water	.	up to 1 fluid ounce

Mix.

MISTURA CRETÆ AROMATICA.

Take of

Prepared Chalk	. .	10 grains
Compound Cinnamon Powder	10 grains
Gum Acacia	. . .	10 grains
Distilled Water	. .	1 fluid ounce

Mix.

MISTURA CRETÆ AROMATICA ANODYNA.

Take of

Aromatic Chalk Mixture	. .	1 fluid ounce
Tincture of Opium	. .	5 minims

Mix.

MISTURA FERRI APERIENS.

Take of

Sulphate of Iron	. .	1 grain
Sulphate of Magnesia	. .	30 grains
Diluted Sulphuric Acid	. .	5 minims
Distilled Water	. .	1 fluid ounce

Mix.

MISTURA FERRI COMPOSITA P.B.

Make according to the instructions given in the British Pharmacopœia, using distilled water in place of rose water.

MISTURA FERRI PERCHLORIDI.

Take of

Tincture of Perchloride of

Iron 10 minims

Glycerine 15 minims

Distilled Water . up to 1 fluid ounce

Mix.

MISTURA FILICIS.

Take of

Liquid Extract of Male Fern 30 minims

Powdered Tragacanth . . 10 grains

Peppermint Water . up to 1 fluid ounce

Rub the liquid extract and the tragacanth together, and then add the peppermint water and mix well.

MISTURA GENTIANÆ CUM AMMONIA.

Take of

Carbonate of Ammonia . 4 grains

Bicarbonate of Soda . . 10 grains

Compound Infusion of

Gentian 1 fluid ounce

Dissolve and mix.

MISTURA GENTIANÆ CATHARTICA.

Take of

Sulphate of Magnesia	.	60 grains
Compound Infusion	of	
Gentian	.	1 fluid ounce

Dissolve.

MISTURA GUAIACI COMPOSITA.

Take of

Ammoniated Tincture of	
Guaiacum	. 30 minims
Mucilage of Gum Acacia	. 1 fluid drachm
Distilled Water	. up to 1 fluid ounce

Mix the tincture with the mucilage;
then add the water and shake well.

MISTURA HYDRARGYRI PERCHLORIDI.

Take of

Perchloride of Mercury	. $\frac{1}{16}$ grain
Tincture of Cochineal	. 10 minims
Distilled Water	. up to 1 fluid ounce

Mix.

MISTURA HYDRARGYRI IODIDI.

Take of

Iodide of Potassium . . .	3 grains
Solution of Perchloride of Mercury	30 minims
Distilled Water	up to 1 fluid ounce

Mix.

MISTURA MAGNESIÆ CUM RHEO.

Take of

Carbonate of Magnesia . . .	15 grains
Rhubarb, in powder	5 grains
Cinnamon Water	1 fluid ounce

Mix.

MISTURA NUCIS VOMICÆ.

Take of

Tincture of Nux Vomica . . .	10 minims
Compound Infusion of Gentian	1 fluid ounce

Mix.

MISTURA POTASSÆ TARTRATIS EFFERVESCENS.

Take of

Bicarbonate of Potash	.	15 grains
Distilled Water	.	4 fluid drachms

Dissolve, and add

Tartaric Acid	.	12 grains
Distilled Water	.	4 fluid drachms

To be taken while effervescing.

MISTURA POTASSII BROMIDI.

Take of

Bromide of Potassium	.	10 grains
Carbonate of Ammonia	.	3 grains
Tincture of Calumba	.	30 minims
Distilled Water	.	1 fluid ounce

Dissolve.

MISTURA POTASSII IODIDI.

Take of

Iodide of Potassium	.	5 grains
Carbonate of Ammonia	.	3 grains
Tincture of Calumba	.	30 minims
Distilled Water	.	up to 1 fluid ounce

Dissolve and mix.

MISTURA QUINIÆ.¹

Take of

Sulphate of Quinia	.	.	1 grain
Dilute Sulphuric Acid	.	.	5 minims
Distilled Water	.	.	1 fluid ounce

Dissolve.

MISTURA QUINIÆ CATHARTICA

Take of

Sulphate of Magnesia	.	.	60 grains
Quinine Mixture	.	.	1 fluid ounce

Dissolve.

MISTURA QUINIÆ CUM FERRO.

Take of

Sulphate of Iron	.	.	1 grain
Quinine Mixture	.	.	1 fluid ounce

Mix.

¹ For out-patients sulphate of cinchonidine and quinine may be substituted for quinia.

MISTURA SCILLÆ COMPOSITA.

Take of

Tincture of Squills . . .	15 minims
Spirit of Nitrous Ether . .	30 minims
Solution of Acetate of Am- monia	1 fluid drachm.
Distilled Water . . . up to	1 fluid ounce

Mix.

MISTURA SENNÆ COMPOSITA.

Take of

Sulphate of Magnesia . . .	120 grains
Tincture of Senna . . .	2 fluid drachms
Compound Tincture of Car- damoms	30 minims
Infusion of Senna . . up to	1 fluid ounce

Dissolve and mix.

PASTA IODI ET AMYLI.

(UNIVERSITY COLLEGE HOSPITAL.)

Take of

Starch, in powder . . .	1 ounce
Glycerine	2 fluid ounces
Water	6 fluid ounces

Boil together, and when nearly cold add

Solution of Iodine B. P. . .	1 fluid ounce
------------------------------	---------------

Mix well.

PILULÆ — PILLS.

PILULA ACONITI ET BELLADONNÆ

Take of

Extract of Aconite . . .	$\frac{1}{6}$ grain
Extract of Belladonna . . .	$\frac{1}{6}$ grain
Sugar of Milk . . .	2 grains
Treacle . . .	a sufficiency

Mix, and make a pill.

PILULA APERIENS.

Take of

Barbadoes Aloes . . .	1 grain
Powdered Scammony . . .	1 grain
Powdered Jalap . . .	1 grain
Powdered Ginger . . .	1 grain
Treacle . . .	a sufficiency

Mix, and make a pill.

**PILULA BELLADONNÆ CUM ZINCI
SULPHATE.**

Take of

Extract of Belladonna . . .	$\frac{1}{8}$ grain
Powdered Sulphate of Zinc . . .	1 grain
Sugar of Milk . . .	1 grain
Treacle . . .	a sufficiency

Mix, and make a pill.

PILULA CALOMELANOS CUM COLO- CYNTHIDE.

Take of

Subchloride of Mercury	. 1 grain
Compound Extract of Co- locynth . . .	3 grains
Extract of Henbane . .	1 grain

Mix, and make a pill.

PILULA CALOMELANOS CUM OPIO.

Take of

Subchloride of Mercury	. 1 grain
Powder of Opium . .	$\frac{1}{4}$ grain
Wheaten Flour . . .	1 grain
Confection of Roses . .	a sufficiency

Mix, and make a pill.

PILULA CALOMELANOS ET PULVERIS IPE- CACUANHÆ COMPOSITI.

Take of

Subchloride of Mercury	. 1 grain
Compound Powder of Ipe- cacuanha . . .	4 grains
Mucilage of Acacia . .	a sufficiency

Mix, and make a pill.

PILULA CANNABIS INDICÆ.

Take of

Extract of Indian Hemp	.	$\frac{1}{4}$ grain
Sugar of Milk	.	2 grains
Treacle	.	a sufficiency

Mix, and make a pill.

PILULA COLCHICI ET QUINIÆ.

Take of

Acetic Extract of Colchi-		
cum	.	1 grain
Sulphate of Quinia	.	1 grain

Mix, and make a pill.

PILULA HYDRARGYRI CUM COLO- CYNTHIDE.

Take of

Mercurial Pill	.	2 grains
Compound Pill of Colo-		
cynth	.	3 grains

Mix, and make a pill.

**PILULA HYDRARGYRI CUM CRETA ET
PULVERIS IPECACUANHÆ COMPOSITI.**

Take of

Mercury with Chalk . . .	2 grains
Compound Powder of Ipe- cacuanha . . .	3 grains
Mucilage of Acacia . . .	a sufficiency

Mix, and make a pill.

PILULA HYDRARGYRI CUM OPIO.

Take of

Mercurial Pill . . .	5 grains
Powder of Opium . . .	$\frac{1}{2}$ grain

Mix, and make a pill.

PILULA HYDRARGYRI CUM RHEO.

Take of

Mercurial Pill . . .	$2\frac{1}{2}$ grains
Compound Rhubarb Pill . .	$2\frac{1}{2}$ grains

Mix, and make a pill.

PILULA SODÆ ARSENIATIS.

Take of

Arseniate of Soda . . .	$\frac{1}{16}$ grain
Sugar of Milk, in powder .	1 grain
Extract of Hop . . .	3 grains

Mix, and make a pill.

PULVERES—POWDERS.

PULVIS ALTERATIVUS.

Take of

Mercury with Chalk .	. 1 part
Bicarbonate of Soda .	. 2 parts
Powdered Rhubarb .	. 3 parts

Mix.

Dose, 6 to 12 grains.

PULVIS CALOMELANOS CUM JALAPA.

Take of

Subchloride of Mercury	. }	each 1 part
Ginger, in powder .	. }	
Jalap, in powder .	. 4 parts	

Mix.

Dose, 12 grains.

PULVIS CALOMELANOS CUM RHEO.

Take of

Subchloride of Mercury	. }	each 1 part
Ginger, in powder .	. }	
Rhubarb, in powder .	. 4 parts	

Mix.

Dose, 12 grains.

PULVIS CALOMELANOS CUM SCAMMONIA.

Take of

Subchloride of Mercury	.	} each 1 part
Ginger, in powder	.	
Scammony, in powder	.	4 parts

Mix.

Dose, 12 grains.

PULVIS CINCHONÆ ET SODÆ.

Take of

Powdered Cinchona Bark	} equal parts
Bicarbonate of Soda	

Mix.

Dose, 5 to 10 grains.

PULVIS HYDRARGYRI CUM CRETA ET BELLADONNÆ.

Take of

Mercury with Chalk	.	2 parts
Powdered Belladonna		
Leaves	.	1 part
Powdered Sugar	.	2 parts

Mix.

Dose, 5 grains.

PULVIS HYDRARGYRI CUM CRETA ET
RHEI.

Take of

Mercury with Chalk .	.	1 part
Rhubarb, in powder .	.	2 parts

Mix.

Dose, 3 to 6 grains.

PULVIS JALAPÆ ET RHEI.

Take of

Jalap, in powder .	.	2 parts
Rhubarb, in powder .	.	1 part

Mix.

Dose, 3 to 15 grains.

PULVIS RHEI ET MAGNESIÆ.

Take of

Rhubarb, in powder .	.	1 part
Carbonate of Magnesia .	.	2 parts

Mix.

Dose, 15 to 20 grains.

PULVIS RHEI ET SODÆ.

Take of

Rhubarb, in powder .	.	} equal parts
Bicarbonate of Soda .	.	

Mix.

Dose, 10 to 20 grains.

PULVIS SENNÆ COMPOSITUS

(VEL PULVIS LIQUIRITIÆ COMPOSITUS PHARM. GERM.).

Take of

Senna Leaves, in powder	. 2 parts
Liquorice Root, in powder	. 2 parts
Fennel Fruit, in powder	. 1 part
Sublimed Sulphur	. 1 part
White Sugar, in powder	. 6 parts

Mix.

Dose, one teaspoonful for an adult.

*UNGUENTA*¹—*OINTMENTS*.**UNGUENTUM BELLADONNÆ COMPOSITUM.**

Take of

Extract of Belladonna	. 80 grains
Distilled Water	. a sufficiency
Ammoniated Mercury, in powder	. 40 grains
Benzoated Lard.	. 360 grains

Rub the extract with water to form a soft paste; then add the ammoniated mercury, and lastly the lard, and mix well.

¹ N.B.—All the ointments may be made with Paraffin Ointment as a basis. This is a mixture of paraffins of a

UNGUENTUM HYDRARGYRI AMMONIATI DILUTUM.

Take of

Ammoniated Mercury, in	
powder	8 grains
Simple Ointment	1 ounce

Mix.

UNGUENTUM HYDRARGYRI AMMONIATI CUM SULPHURE.

Take of

Ammoniated Mercury, in	
powder	6 grains
Sublimed Sulphur	30 grains
Benzoated Lard	1 ounce

Mix.

UNGUENTUM HYDRARGYRI NITRATIS MITE.

Take of

Ointment of Nitrate of	
Mercury	1 part
Paraffin Ointment	7 parts

Rub them well together.

soft unctuous consistency, having a melting point of 95° to 100° F. It is odourless, has a bland taste, and may be of a white opal or pale yellow colour. It has a minute crystalline structure not visible to the naked eye. It does not become rancid, and is not affected by boiling with a caustic alkaline solution. In commerce the substances known as Vaseline and Cosmoline (unguentum petrolei) may be taken as representing it.

UNGUENTUM HYDRARGYRI NITRATIS MITIUS.

Take of

Ointment of Nitrate of

Mercury . . . 1 part

Paraffin Ointment . . 11 parts

Mix.

UNGUENTUM HYDRARGYRI OXIDI FLAVI.

Take of

Yellow Oxide of Mercury . 8 grains

Paraffin Ointment . . 1 ounce

Mix.

UNGUENTUM HYDRARGYRI OXIDI RUBRI DILUTUM.

Take of

Ointment of Red Oxide of

Mercury . . . 1 part

Paraffin Ointment . . 7 parts

Rub them well together.

UNGUENTUM HYDRARGYRI CUM BELLA- DONNA.

Take of

Extract of Belladonna . 2 parts

Distilled Water . . a sufficiency

Ointment of Mercury . 6 parts

Rub the extract with water to form a soft paste; add the ointment, and mix well.

UNGUENTUM PLUMBI SUBACETATIS CUM PARAFFINE.

Take of

Glycerole of Subacetate of

Lead	4½ fluid ounces
Paraffin Ointment . .	18 ounces
Paraffin Wax	6 ounces

Melt the paraffin wax and ointment together, add the glycerole of lead, and stir till cold.

UNGUENTUM SIMPLEX.

Take of

White Wax	1 part
Prepared Lard	2 parts

Melt them together.

UNGUENTUM STAPHISAGRIÆ.

Take of

Stavesacre, in powder .	1 part
Olive Oil	1 part
Prepared Lard	2 parts

Melt the lard in a water bath, add the oil and stavesacre, and maintain the heat for a quarter of an hour, then remove and stir till cold.

UNGUENTUM ZINCI COMPOSITUM.

Take of

Ointment of Zinc	.		} each equal parts
Compound Ointment of			
Subacetate of Lead B.P.			

Rub them well together.

UNGUENTUM ZINCI OLEATIS.

Take of


Oxide of Zinc	.	.	.	1 ounce
Oleic Acid	.	.	.	8 fluid ounces


Mix, and set aside for six hours; then apply heat to liquefy and form a perfect solution.

To this add

Paraffin Ointment	.	.	.	9 ounces
-------------------	---	---	---	----------

Dissolve, and stir till cold.





BOUND BY
POTTISWOODE & CO.
LONDON.

